

RENEWAL SERIES AND SQUARE-ROOT BOUNDARIES FOR BESSEL PROCESSES

NATHANAËL ENRIQUEZ, CHRISTOPHE SABOT, AND MARC YOR

Abstract. We show how a description of Brownian exponential functionals as a renewal series gives access to the law of the hitting time of a square-root boundary by a Bessel process. This extends classical results by Breiman and Shepp, concerning Brownian motion, and recovers by different means, extensions for Bessel processes, obtained independently by Delong and Yor.

Let B_t be the standard real valued Brownian motion and for $\nu > 0$, introduce the geometric Brownian motion $\mathcal{E}_t^{(-\nu)}$ and its exponential functional $\mathcal{A}_t^{(-\nu)}$

$$\begin{aligned}\mathcal{E}_t^{(-\nu)} &:= \exp(B_t - \nu t) \\ \mathcal{A}_t^{(-\nu)} &:= \int_0^t (\mathcal{E}_s^{(-\nu)})^2 ds.\end{aligned}$$

Lamperti's representation theorem [5] applied to $\mathcal{E}_t^{(-\nu)}$ states

$$(0.1) \quad \mathcal{E}_t^{(-\nu)} = R_{\mathcal{A}_t^{(-\nu)}}^{(-\nu)}$$

where $(R_u^{(-\nu)}, u \leq T_0(R^{(-\nu)}))$ denotes the Bessel process of index $(-\nu)$ (equivalently of dimension $\delta = 2(1 - \nu)$), starting at 1, which is an $\mathbb{R}_+$ -valued diffusion with infinitesimal generator $\mathcal{L}^{(-\nu)}$ given by

$$\mathcal{L}^{(-\nu)} f(x) = \frac{1}{2} f''(x) + \frac{1 - 2\nu}{2x} f'(x), \quad f \in C_b^2(\mathbb{R}_+^*).$$

Let us remark that, in the special case $\nu = 1/2$, equation (0.1) is nothing else but the Dubins-Schwarz representation of the exponential martingale $\mathcal{E}_t^{(-1/2)}$ as Brownian motion time changed with $\mathcal{A}_t^{(-1/2)}$.

For a short summary of relations between Bessel processes and exponentials of Brownian motion, see e.g. Yor [10].

Let us consider now the following random variable Z , which is often called a perpetuity in the mathematical finance literature:

$$Z := \mathcal{A}_\infty^{(-\nu)} = \int_0^\infty (\mathcal{E}_s^{(-\nu)})^2 ds$$

We deduce directly from (0.1) that

$$\mathcal{A}_\infty^{(-\nu)} = T_0(R^{(-\nu)})$$

2000 *Mathematics Subject Classification.* 60G40, 60J57.

Key words and phrases. Bessel processes, renewal series, exponential functionals, square-root boundaries.

where $T_0 := \inf\{u : X_u = 0\}$, and it is well-known (see [4], [11]), that

$$(0.2) \quad \mathcal{A}_\infty^{(-\nu)} \stackrel{(law)}{=} \frac{1}{2\gamma_\nu}$$

where γ_ν is a gamma variable with parameter ν (i.e. with density $\frac{1}{\Gamma(\nu)}x^{\nu-1}e^{-x}\mathbf{1}_{\mathbb{R}_+}$).

Our main result characterizes the law of the hitting time of a parabolic boundary by $R_u^{(-\nu)}$ which corresponds to a Bessel process of dimension $d < 2$.

Theorem 1. *Let $0 < b < c$, and $\sigma := \inf\{u : (R_u^{(-\nu)})^2 = \frac{1}{c}(b + u)\}$ with $R_0^{(-\nu)} = 1$.*

$$(0.3) \quad E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(1 + 2b\gamma_{\nu+s})^{-s}]}{E[(1 + 2c\gamma_{\nu+s})^{-s}]}, \quad \text{for any } s \geq 0$$

Proof: using the strong Markov property and the stationarity of the increments of Brownian motion, we obtain that for any stopping time τ of the Brownian motion

$$\mathcal{A}_\infty^{(-\nu)} =: Z = \mathcal{A}_\tau^{(-\nu)} + (\mathcal{E}_\tau^{(-\nu)})^2 Z'$$

where Z' is independent of $(\mathcal{A}_\tau^{(-\nu)}, \mathcal{E}_\tau^{(-\nu)})$ and $Z \stackrel{(law)}{=} Z'$.

This implies, by (0.1), that Z satisfies the following affine equation (see [8] for a survey about these equations)

$$(0.4) \quad \mathcal{A}_\infty^{(-\nu)} =: Z = \mathcal{A}_\tau^{(-\nu)} + (R_{\mathcal{A}_\tau^{(-\nu)}}^{(-\nu)})^2 Z'$$

where Z' is independent of $(\mathcal{A}_\tau^{(-\nu)}, R_{\mathcal{A}_\tau^{(-\nu)}}^{(-\nu)})$ and $Z \stackrel{(law)}{=} Z'$.

Obviously, $\sigma < T_0(R^{(-\nu)})$. Taking now :

$$\tau = \inf\{t : (R_{\mathcal{A}_t^{(-\nu)}}^{(-\nu)})^2 = \frac{1}{c}(b + \mathcal{A}_t^{(-\nu)})\}$$

we get $\mathcal{A}_\tau^{(-\nu)} = \sigma$, and the identity in law

$$(0.5) \quad b + Z \stackrel{(law)}{=} (b + \sigma)(1 + \frac{Z}{c})$$

where the variables σ and Z on the right-hand side are independent.

As a result, we obtain the Mellin-Stieltjes transform of σ :

$$E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(b + Z)^{-s}]}{E[(c + Z)^{-s}]}$$

But, from (0.2)

$$E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(2\gamma_\nu)^s \frac{1}{(1+2b\gamma_\nu)^s}]}{E[(2\gamma_\nu)^s \frac{1}{(1+2c\gamma_\nu)^s}]}$$

which gives the result. $\square$

One can now use the duality between the laws of Bessel processes of dimension d and $4 - d$ to get the analogous result of Theorem 1, and recover the result of Delong [2], [3], and Yor [9] which deals with the case $d \geq 2$.

Theorem 2. *Let $0 < b < c$, and $\sigma := \inf\{u : (R_u^{(\nu)})^2 = \frac{1}{c}(b + u)\}$ with $R_0^{(\nu)} = 1$.*

$$(0.6) \quad E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(1 + 2b\gamma_s)^{-s+\nu}]}{E[(1 + 2c\gamma_s)^{-s+\nu}]}, \quad \text{for any } s \geq 0.$$

Proof : it is based on the following classical relation between the laws of the Bessel processes with indices ν and $-\nu$:

$$(0.7) \quad \mathcal{P}_x^{(\nu)}|_{\mathcal{F}_t} = \frac{(X_{t \wedge T_0})^{2\nu}}{x^{2\nu}} \cdot \mathcal{P}_x^{(-\nu)}|_{\mathcal{F}_t}$$

which implies that

$$E_1^{(\nu)}[(b + \sigma)^{-s}] = E_1^{(-\nu)}[X_\sigma^{2\nu}(b + \sigma)^{-s}] = \frac{1}{c^\nu} E_1^{(-\nu)}[(b + \sigma)^{-s+\nu}]$$

Theorem 1 gives the result. $\square$

Finally, it is easily shown, thanks to the classical representations of the Whittaker functions (see Lebedev [6]), that the right-hand sides of (0.3) and (0.6) are expressed in terms of ratios of Whittaker functions.

Acknowledgement: We would like to thank Daniel Dufresne for useful and enjoyable discussions on the subject.

REFERENCES

- [1] Breiman, L. First exit times from a square root boundary. (1967) Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. II: Contributions to Probability Theory, Part 2 pp. 9–16 Univ. California Press, Berkeley, Calif.
- [2] Delong, D.M. Crossing probabilities for a square-root boundary for a Bessel process. Comm. Stat. A - Theory Methods. 10 (1981), no 21, 2197–2213.
- [3] Delong, D. M. Erratum: "Crossing probabilities for a square root boundary by a Bessel process" Comm. Statist. A - Theory Methods 12 (1983), no. 14, 1699.
- [4] Dufresne, D. The distribution of a perpetuity, with applications to risk theory and pension funding. Scand. Actuarial J. (1990) 39–79.
- [5] Lamperti, J. Semi-stable Markov processes. I. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 22 (1972), 205–225.
- [6] Lebedev, N. N. Special functions and their applications. Dover (1972).
- [7] Shepp, L. A. A first passage problem for the Wiener process. Ann. Math. Statist. 38 (1967) 1912–1914.
- [8] Vervaat, W. On a stochastic difference equation and a representation of nonnegative infinitely divisible random variables. Adv. in Appl. Probab. 11 (1979), no. 4, 750–783.
- [9] Yor, M. On square-root boundaries for Bessel processes, and pole-seeking Brownian motion. Stochastic analysis and applications (Swansea, 1983), 100–107, Lecture Notes in Math., 1095, Springer, Berlin, 1984.
- [10] Yor, M. On some exponential functionals of Brownian motion. Adv. Appl. Prob. 24 (1992), 509–531.
- [11] Yor, M. Sur certaines fonctionnelles exponentielles du mouvement brownien réel. J. Appl. Prob 29 (1992), 202–208.

LABORATOIRE MODAL'X, UNIVERSITÉ PARIS 10, 200 AVENUE DE LA RÉPUBLIQUE, 92000
NANTERRE, FRANCE

E-mail address: `nenriquez@u-paris10.fr`

UNIVERSITÉ DE LYON, UNIVERSITÉ LYON 1, INSTITUT CAMILLE JORDAN, CNRS UMR 5208,
43, BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE

E-mail address: `sabot@math.univ-lyon1.fr`

LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, CNRS UMR 7599, UNIVERSITÉ
PARIS 6, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address: `deaproba@proba.jussieu.fr`